

# Standard and non-standard neutrino properties

**Antonio Palazzo**



MAX-PLANCK-GESELLSCHAFT



MARIE CURIE  
ACTIONS



SEVENTH FRAMEWORK  
PROGRAMME

**Max-Planck-Institut für Physik (Munich)**

# Outline

- 1) The path to the discovery of non-zero  $\theta_{13}$
- 2) The hints of sterile neutrinos and the Sun
- 3) Solar  $\nu$ s as a probe of the MSW dynamics

Tightly interconnected topics, as we will see ...

# The path to the discovery of non-zero $\theta_{13}$

# Why a non-zero $\theta_{13}$ is so important

$$J = \Im[U_{\mu 3} U_{e 2} U_{\mu 2}^* U_{e 3}^*]$$

The Jarlskog invariant  $J$  gives a parameterization-independent measure of the CP violation induced by the complexity of  $U$

In the standard parameterization the expression of  $J$  is:

$$J = \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta$$

Only if all three  $\theta_{ij} \neq 0$  we can have CP violation

quark-sector:  $J_{\text{CKM}} \sim 3 \times 10^{-5}$ , much smaller than  $|J|_{\text{max}} = \frac{1}{6\sqrt{3}} \sim 0.1$

lepton-sector:  $|J|$  may be as large as  $3 \times 10^{-2}$  (it will depend on  $\delta$ )

# Historical result established by CHOOZ in 1998

$$P_{ee}^{\text{osc}} = 1 - 4U_{e3}^2(1 - U_{e3}^2) \sin^2 \left( \frac{\Delta m^2}{4E} L \right)$$


$$P_{ee}^{\text{exp}} \simeq 1 \quad U_{e3}^2 = \sin^2 \theta_{13}$$



Exclusion plot in the  $(\Delta m^2, \theta_{13})$  plane

$\Delta m^2$  scale now set with precision by **Atm +LBL**

CHOOZ exclusion plot



# ...since then...


The  $3\nu$  global analyses have played an increasingly relevant role in pinning down  $\theta_{13}$ , constantly improving their sensitivity.

They have first corroborated (atm. analyses) and then strengthened (sol+Kam analyses) the CHOOZ upper limit.

Hence, in 2008 it was not surprising that they started to be competitive, reaching values of  $\theta_{13}$  below the CHOOZ limit.

What instead - pleasantly - surprised us was that, for the first time, the analyses pointed towards a non-zero value of this parameter...

# 2008: Global 3ν analysis




preference for  $\theta_{13} > 0$

The global analysis provided a preference for  $\theta_{13} > 0$  at 90% C.L.

Fogli, Lisi, Marrone, A.P., Rotunno,  
PRL 101, 141801 (2008), arXiv:0806.2649, hep-ph.


# 2008: First indication of non-zero $\theta_{13}$



Fogli, Lisi, Marrone, A.P., Rotunno, Phys. Rev. Lett. 101, 141201 (2008)

Two independent hints came from solar and atmospheric sectors:


$$\sin^2\theta_{13} \sim 0.016$$






# Indication irrefutably confirmed in 2012


## Daya Bay



## Reno




## Double-CHOOZ



# Parameter estimates as of June 2012

Synopsis of global 3ν oscillation analysis



$\theta_{13}$  non-zero at  $8\sigma$


$\sim 2\sigma$  indication of non-maximal  $\theta_{23}$  ( $\theta_{23} < \pi/4$ )

Hint of  $\delta \sim \pi$

no sensitivity to mass hierarchy

Fogli, Iasi, Marrone, Montanino, A.P., Rotunno, PRD 86 013012 (2012)  
(includes Neutrino 2012 results)

# How the indication of $\theta_{23} < \pi/4$ emerges



- LBL introduce:**
- $\theta_{23}$ - $\theta_{13}$  anticorrelation
  - prefer. non-maximal  $\theta_{23}$
  - weak octant asymmetry


Once reactors fix  $\theta_{13}$   
the octant asymmetry  
is enhanced

Atm. further enhance  
octant asymmetry

**Global indication of  
 $\theta_{23} < \pi/4$  emerges**

Fogli, Iasi, Marrone, Montanino, A.P., Rotunno, PRD 86 013012 (2012)  
(includes Neutrino 2012 results)

# First information about $\delta$



LBL are almost insensitive to  $\delta$

Weak sensitivity emerges once reactors fix  $\theta_{13}$

Atm. enhance sensitivity

Global hint of  $\delta \sim \pi$  emerges


Fogli, Iliu, Marrone, Montanino, A.P., Rotunno, PRD 86 013012 (2012)  
(includes Neutrino 2012 results)

If  $\delta \sim \pi$  confirmed it would indicate  $U \sim$  real and a small  $J$   
... and a long and difficult way towards CPV observation!

But let us come back to the solar hint of  $\theta_{13} > 0$

A closer look shows that it emerged from a delicate interplay of solar and KamLAND

$\theta_{13} = 0$



$\theta_{13} > 0$

Solar  $\nu$ s are thus a very precise machine and we can trust it also when searching for non-standard physics!

**Beyond the standard  $3\nu$  paradigm**

**Exploring new neutrino properties**

# Why go beyond the standard 3 $\nu$ picture?

## Theory

Many extensions of the SM point towards new  $\nu$  properties (interactions, new states,...)

## Acquired knowledge

Precision on standard parameters enhances the sensitivity to any kind of perturbation

## Experimental hints

Although the 3 $\nu$  scheme explains most of the data an increasing number of anomalies is showing up


## Future data

A rich plan of new experiments will allow us to explore and chart unknown territories

# The hints of sterile neutrinos and the Sun



# Hint #1: The Gallium calibration anomaly




SAGE coll., PRC 73 (2006) 045805


**Deficit observed in calibration performed with radioactive sources**

**But it could be due to overestimate of  $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$  cross section**

# Hint #2: The reactor antineutrino anomaly



Mention et al., PRD 83 073006 (2011)




Mueller et al., PRC 83 054615 (2011)  
Huber, PRC 84 024617 (2011)

With new reactor fluxes deficit of all the short-baseline reactor measurements

But new calculations, like older ones, are still anchored to (one single)  $\beta$ -spectrum experiment (ILL)


# Fitting the short-distance $\nu_e$ -disappearance




Mention et al., PRD 83 073006 (2011)

$$\sin^2 2\theta_{new} \simeq 0.1 \quad \Delta m_{new}^2 \gtrsim 1 \text{ eV}^2$$

# Hint #3: Anomalous short-distance $\nu_e$ -appearance



LSND, PRL 75 (1995) 2650



Giunti and Laveder, arXiv:1107.1452

## Warning:


In tension with disappearance searches:  
 $\nu_\mu \rightarrow \nu_e$  positive appearance signal incompatible with  
 joint  $\nu_e \rightarrow \nu_e$  (positive) &  $\nu_\mu \rightarrow \nu_\mu$  (negative) searches

Theory:

$$\sin^2 2\theta_{e\mu} \simeq \frac{1}{4} \sin^2 2\theta_{ee} \sin^2 2\theta_{\mu\mu} \simeq 4|U_{e4}|^2 |U_{\mu4}|^2$$

Experiments:  $\sim$  few %       $\sim$  0.1       $<$  few %

## Hint #4: Cosmology favors extra radiation



CMB + LSS tend to prefer  
extra relativistic content  
~ 2 sigma effect

[Hamann et al., PRL 105, 181301 (2010)]


### Warnings:

- **eV masses acceptable only abandoning standard  $\Lambda$ CDM**  
(Kristiansen & Elgaroy arXiv:1104.0704 , Hamann et al. arXiv:1108.4136)
- **$N_s > 1$  at BBN strongly disfavored** (Mangano & Serpico PLB 701, 296, 2011)
- **$N_s$  is not specific of  $\nu_s$**   
(new light particles, decay of dark matter particles, quintessence, ...)

Can we get some information on  $\nu_s$   
from the solar neutrino sector?


# The 3+1 scheme:

The 4<sup>th</sup>  $\nu$  state induces a small perturbation of the 3-flavor framework




$$\Delta m_{\text{atm}}^2$$

$$\Delta m_{\text{sol}}^2$$




$$|U_{s4}| \sim 1$$




$$\Delta m_{\text{new}}^2 > 1 \text{eV}^2$$

From the "point of view" of the solar doublet ( $\nu_1, \nu_2$ ) we expect similar sensitivity to  $U_{e3}$  &  $U_{e4}$

# VSBL $\nu_e$ disappearance in a 3+1 scheme



Mention et al. arXiv:1101:2755 [hep-ex]



SAGE coll., PRC 73 (2006) 045805

In a 2ν framework:

$$P_{ee} \simeq 1 - \sin^2 2\theta_{new} \sin^2 \frac{\Delta m_{new}^2 L}{4E}$$

In a 3+1 scheme:

$$P_{ee} = 1 - 4 \sum_{j>k} U_{ej}^2 U_{ek}^2 \sin^2 \frac{\Delta m_{jk}^2 L}{4E}$$

$$\Delta m_{sol}^2 \ll \Delta m_{atm}^2 \ll \Delta m_{new}^2$$

$$\sin^2 \theta_{new} \simeq U_{e4}^2 = \sin^2 \theta_{14}$$

3+1 scheme has several consequences: solar, atm, react., accel.

We will focus on the implications for Solar (S) & KamLAND (K)



# LBL $\nu_e$ disappearance in a 3+1 scheme

$$P_{ee} = 1 - 4 \sum_{j>k} U_{ej}^2 U_{ek}^2 \sin^2 \frac{\Delta m_{jk}^2 L}{4E}$$


$$\Delta m_{sol}^2 \ll \Delta m_{atm}^2 \ll \Delta m_{new}^2$$

$\Delta m_{atm}^2$   
 $\Delta m_{new}^2$  -driven osc. averaged

$$P_{ee} = (1 - U_{e3}^2 - U_{e4}^2)^2 P_{ee}^{2\nu} + U_{e3}^4 + U_{e4}^4$$

$$U_{e3}^2 = c_{14}^2 s_{13}^2 \quad U_{e4}^2 = s_{14}^2$$

KamLAND



Exact degeneracy between  $U_{e3}$  and  $U_{e4}$

# Solar $\nu$ conversion in a 3+1 scheme

$$i \frac{d}{dx} \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ \nu_s \end{pmatrix} = H \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \\ \nu_s \end{pmatrix} \quad H = UKU^T + V(x)$$

$$K = \frac{1}{2E} \text{diag}(k_1, k_2, k_3, k_4) \quad k_i = \frac{m_i^2}{2E} \quad \text{wavenumbers in vacuum}$$

Useful to write the mixing matrix as\*:  $U = R_{23} \mathbf{S} R_{13} R_{12} \quad \mathbf{S} = R_{24} R_{34} R_{14}$

$$\theta_{14} = \theta_{24} = \theta_{34} = 0 \quad \text{-->} \quad \mathbf{S} = \mathbf{I} \quad \text{-->} \quad \text{3-flavor case}$$

$$V = \text{diag}(V_{CC}, 0, 0, -V_{NC}) \quad \text{MSW potential}$$

$$V_{CC} = \sqrt{2} G_F N_e \quad V_{NC} = \frac{1}{2} \sqrt{2} G_F N_n$$

\* We assume  $U$  to be real but in general it can be complex due to CP phases

**Change of basis:**  $\nu' = (R_{23} S R_{13})^T \nu = A^T \nu = R_{12} U^T$

**In the new basis:**  $H' = A^T H A = R_{12} K R_{12}^T + R_{13}^T S^T V S R_{13}$

**At zero<sup>th</sup> order in:**

$\frac{V}{k_{atm}}$  and  $\frac{V}{k_{new}}$

$$H' \simeq \begin{pmatrix} H'_{2\nu} & & & \\ \text{---} & & & \\ & & k_3 & \\ & & & k_4 \\ & & & & \vdots \end{pmatrix}$$

The 3<sup>rd</sup> & 4<sup>th</sup> state evolve independently from the 1<sup>st</sup> & 2<sup>nd</sup>

The dynamics reduces to that of a 2x2 system

# Diagonalization of the Hamiltonian

The 2x2 Hamiltonian  
is diagonalized by a  
1-2 rotation

$$\tilde{R}_{12}^T H'_{2\nu} \tilde{R}_{12} = \text{diag}(\tilde{k}_1, \tilde{k}_2)$$

which defines the solar  
mixing angle in matter

$$\tilde{\theta}_{12}(x)$$

wavenumbers in matter

$$\tilde{k}_i$$

The starting Hamiltonian  
is then diagonalized by

$$\tilde{U} = A \tilde{R}_{12}$$

$$\tilde{U}^T H \tilde{U} = \text{diag}(\tilde{k}_1, \tilde{k}_2, k_3, k_4)$$

For  $\nu_3$  and  $\nu_4$  (averaged) vacuum-like propagation

**The 2x2 Hamiltonian:**  $H'_{2\nu} = H'_{2\nu}{}^{\text{kin}} + H'_{2\nu}{}^{\text{dyn}}$

$$H'_{2\nu}{}^{\text{kin}} = \begin{pmatrix} c_{12} & s_{12} \\ -s_{12} & c_{12} \end{pmatrix} \begin{pmatrix} -k_{\text{sol}}/2 & 0 \\ 0 & k_{\text{sol}}/2 \end{pmatrix} \begin{pmatrix} c_{12} & -s_{12} \\ s_{12} & c_{12} \end{pmatrix} \quad k_{\text{sol}} = \frac{m_2^2 - m_1^2}{2E}$$

$$H'_{2\nu}{}^{\text{dyn}} = V_{CC}(x) \begin{pmatrix} \gamma^2 + r(x) \alpha^2 & r(x) \alpha \beta \\ r(x) \alpha \beta & r(x) \beta^2 \end{pmatrix}^* \quad \text{Formally equivalent to NSI (see later)}$$

$$\begin{cases} \alpha^2 + \beta^2 = U_{s1}^2 + U_{s2}^2 \\ \gamma^2 = 1 - (U_{e3}^2 + U_{e4}^2) \end{cases} \quad \begin{cases} \alpha = c_{24}c_{34}c_{13}s_{14} - s_{34}s_{13} \\ \beta = s_{24}c_{34} \\ \gamma = c_{13}c_{14} \end{cases} \quad r(x) = \frac{V_{NC}(x)}{V_{CC}(x)}$$

**New MSW dynamical corrections induced by the 4<sup>th</sup> state are smaller than 1% and too small to be observable (see later).**

**But important new kinematical effects are present ...**

For adiabatic propagation (valid for small deviations around the LMA)

$$P_{ee} = \sum_{i=1}^4 U_{ei}^2 \tilde{U}_{ei}^2 = U_{e1}^2 \tilde{U}_{e1}^2 + U_{e2}^2 \tilde{U}_{e2}^2 + U_{e3}^4 + U_{e4}^4$$

$$P_{es} = \sum_{i=1}^4 U_{si}^2 \tilde{U}_{ei}^2 = U_{s1}^2 \tilde{U}_{e1}^2 + U_{s2}^2 \tilde{U}_{e2}^2 + U_{s3}^2 U_{e3}^2 + U_{s4}^2 U_{e4}^2$$

Expressions for  $U_{ei}$ 's  
(always valid)

$$\left. \begin{aligned} U_{e1}^2 &= c_{14}^2 c_{13}^2 c_{12}^2 \\ U_{e2}^2 &= c_{14}^2 c_{13}^2 s_{12}^2 \\ U_{e3}^2 &= c_{14}^2 s_{13}^2 \sim s_{13}^2 \\ U_{e4}^2 &= s_{14}^2 \end{aligned} \right\} \sim 1 - s_{14}^2 - s_{13}^2$$

Expressions for  $U_{si}$ 's  
valid for  $\theta_{24} = \theta_{34} = 0$

$$\left. \begin{aligned} U_{s1}^2 &= s_{14}^2 c_{13}^2 c_{12}^2 \\ U_{s2}^2 &= s_{14}^2 c_{13}^2 s_{12}^2 \\ U_{s3}^2 &= s_{14}^2 s_{13}^2 \sim 0 \\ U_{s4}^2 &= c_{14}^2 c_{13}^2 \sim 1 - s_{14}^2 \end{aligned} \right\} \sim s_{14}^2$$

The elements of  $\tilde{U}$  are obtained replacing  $\theta_{12}$  with  $\tilde{\theta}_{12}$  calculated in the production point (near the sun center)



# Solar $\nu$ : Two simple limit cases

$$\theta_{13} \neq 0 \quad \theta_{14} = 0 \quad (3\nu)$$


$$\begin{cases} P_{ee} = c_{13}^4 P_{ee}^{2\nu} \Big|_{V \rightarrow Vc_{13}^2} + s_{13}^4 \\ P_{es} = 0 \end{cases}$$

$$\theta_{13} = 0 \quad \theta_{14} \neq 0 \quad (4\nu)$$

$$\begin{cases} P_{ee} = c_{14}^4 P_{ee}^{2\nu} \Big|_{V \rightarrow Vc_{14}^2} + s_{14}^4 \\ P_{es} \simeq s_{14}^2 P_{ee}^{2\nu} \Big|_{V \rightarrow Vc_{14}^2} + s_{14}^2 \end{cases}$$




# $(\theta_{13}, \theta_{12})$ vs $(\theta_{14}, \theta_{12})$ constraints



$$\begin{cases} CC \sim \Phi_B P_{ee} \\ NC \sim \Phi_B (1 - P_{es}) \\ ES \sim \Phi_B (P_{ee} + 0.15 P_{ea}) \end{cases}$$

Solar  $\nu$  sensitive to  $P_{es}$   
 CC/NC (SNO) & ES (SK)




But unfortunately only small differences among  $3\nu$  and  $4\nu$

We expect a degeneracy among  $\theta_{13}$  and  $\theta_{14}$



# $(\theta_{13}, \theta_{14})$ constraints



**Complete degeneracy**  
 $\theta_{13}-\theta_{14}$  indistinguishable

**Solar sector essentially sensitive to  $\sim U_{e3}^2 + U_{e4}^2$**


**Hint for  $\nu_e$  mixing with states others than  $(\nu_1, \nu_2)$**

**Different probes are necessary to determine if  $\nu_e$  mixes with  $\nu_3$  or  $\nu_4$**

A.P. PRD 83 113013 (2011) [arXiv: 1105.1705 hep-ph]

# Evidence of $\theta_{13} > 0$ kills preference of $\theta_{14} > 0$


UPDATE of: A.P. PRD 85 077302 (2012) [arXiv: 1201.4280 hep-ph]



- Upper limit  $\rightarrow \sin^2 \theta_{14} < 0.04$  (90% C.L.)
- KamLAND, only spectral shape included:  
limit is independent of reactor flux estimates
- $\theta_{13}$  estimate independent of  $\theta_{14}$

Solar bound is the most stringent one for  $\Delta m_{14}^2 > 1 \text{eV}^2$

Compilation of all the existing limits on  $\theta_{14}$  ( $= \theta_{ee}$ )




Talk by C. Giunti @  $\nu$ TURN 2012

# How to go below the solar upper limit

Make use of a  $\nu$  source close to a Borexino-like detector

Talk by M. Pallavicini @ Neutrino 2012



Identified by reactor anomaly

Solar upper limit

Borexino sensitivity

## Kairos (καιρός)




### The "Borexino Kairos"

This is the right (and fleeting) moment (kairos) for Borexino to exploit its unique potential!

# Solar $\nu$ s as a probe of non-standard MSW dynamics

# Coherent forward scattering in the presence of NSI : pictorial view



NSI described  
by an effective  
four-fermions  
operator

$$O_{\alpha\beta}^{\text{NSI}} \sim \bar{\nu}_\alpha \nu_\beta \bar{f} f$$

$$(\alpha, \beta) = e, \mu, \tau$$

$$f \equiv (e, u, d)$$

# Coherent forward scattering in the presence of NSI : math. view

**Evolution in the flavor basis:** 
$$i \frac{d}{dx} \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = H \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix}$$

**H contains three terms:** 
$$H = H_{\text{kin}} + H_{\text{dyn}}^{\text{std}} + H_{\text{dyn}}^{\text{NSI}}$$

**Kinematics** 
$$H_{\text{kin}} = U \begin{pmatrix} -\delta k/2 & 0 & 0 \\ 0 & +\delta k/2 & 0 \\ 0 & 0 & k/2 \end{pmatrix} U^\dagger \quad \begin{aligned} \delta k &= \delta m^2 / 2E \\ k &= m^2 / 2E \end{aligned}$$

**Standard  
MSW  
dynamics**

$$H_{\text{dyn}}^{\text{std}} = \text{diag}(V, 0, 0) \quad V(x) = \sqrt{2} G_F N_e(x)$$

**Non-standard  
dynamics**

$$(H_{\text{dyn}}^{\text{NSI}})_{\alpha\beta} = \sqrt{2} G_F N_f(x) \epsilon_{\alpha\beta}$$

# Reduction to an effective two flavor dynamics

One mass scale approximation:  $\Delta m^2 \rightarrow \infty$

$$P_{ee} = c_{13}^4 P_{ee}^{\text{eff}} + s_{13}^4 \quad \text{survival probability}$$

$$i \frac{d}{dx} \begin{pmatrix} \nu_e \\ \nu_a \end{pmatrix} = H^{\text{eff}} \begin{pmatrix} \nu_e \\ \nu_a \end{pmatrix} \quad \text{effective evolution}$$

$$H^{\text{eff}} = V(x) \begin{pmatrix} c_{13}^2 & 0 \\ 0 & 0 \end{pmatrix} + \sqrt{2} G_f N_d(x) \begin{pmatrix} 0 & \varepsilon \\ \varepsilon & \varepsilon' \end{pmatrix} \quad \text{Formally similar to } 4\nu \text{ effects}$$

For  $\theta_{13} = 0$  :

$$\varepsilon = -\varepsilon_{e\mu} c_{23} - \varepsilon_{e\tau} s_{23}$$

$$\varepsilon' = -2\varepsilon_{\mu\tau} s_{23} c_{23}$$


$\varepsilon_{\mu\tau} \sim 0$  (strong bounds from atmospheric  $\nu$ )

Parameter space:

$$[\delta m^2, \theta_{12}, \varepsilon]$$



# Impact of NSI on the solar spectrum



$$\epsilon = -0.16$$


$$(\epsilon_{e\tau} = +0.23)$$

for interaction  
with d-quark


NSI with a size of  $\sim 10\%$  are needed to produce appreciable effects:  
 $4\nu$  effects induced by sterile neutrinos ( $\sim 1\%$ ) are thus unobservable

# NSIs can help to explain the anomalous spectrum behavior


## BOREXINO



## SNO



## SK



# This hypothesis can be tested quantitatively

The response functions of SK, SNO, Borexino are centered around  $E_0 = 10$  MeV, where they have maximal sensitivity


Assuming a regular behavior for the survival probability we can parameterize its high energy behavior as a second order polynomial

$$P_{ee} = c_0 + c_1 (E-E_0) + c_2 (E-E_0)^2$$

It is then possible to:

- 1) Extract the coefficients from the combination of all the experiments sensitive to the high-energy neutrinos.
- 2) Check where a given theor. model (standard MSW, +NSI, etc.) “lives” in the space of the coefficients  $c_i$ ’s.


# Constraints on $[c_1, c_2]$




A.P, PRD 83, 101701 (2011) (Rapid Communications) arXiv:1101.3875

NSI gains a  $\Delta\chi^2 \sim -2.0$  from better description of the spectrum

# NSI can also alleviate tension in $\delta m^2$ determinations



M. Smy @ Neutrino 2012



A.P. and J.W.F. Valle, PRD 80, 091301 (2009)

---

---

# Summary

---

---

- $3\nu$  paradigm acquires a new crucial piece:  $\theta_{13} > 0$
- First interesting information on CPV phase ( $\delta \sim \pi$ ) &  $\theta_{23} < \pi/4$
- A few expts. results suggest  $\nu$ 's mix with new sterile states
- Evidence of  $\theta_{13} > 0$  + solar sector data provide the stringent and robust upper limit:  $U_{e4}^2 < 0.04$  (90% C.L.)
- The solar sector data evidence two weak anomalies which can be explained in terms of new neutrino interactions.
- New experiments indispensable to settle the issues.

**Thank you  
for  
your attention!**